



New Approach of Non-Linear Fractional Differential Equations Analytical Solution by Akbari-Ganji's Method

Ziba Khalili¹, Ali Bararpour¹, Khashayar Hosseinzadeh^{1*}, Bahram Jafari², Davood Domiri Ganji¹

¹ Department of Mechanical Engineering, Babol Noshirvani University of Technology, Babol, Iran.

² Faculty of Engineering Modern Technologies, Amol University of Special Modern Technologies (AUSMT), Amol, Iran.

Article Info

Received 05 January 2024

Accepted 10 February 2024

Available online 15 March 2024

Keywords:

Fractional Differential Equations;

Akbari-Ganji's Method;

Mathematical Methods.

Abstract:

The present study examines fractional differential equations (FDEs) as these types of equations are widely used in modeling. FDEs are more difficult to solve than differential equations, which have integer order. There are a variety of mathematical methods for solving these equations. An analytical method is utilized in this paper to solve non-linear FDEs. Akbari-Ganji's method (AGM) has been used as a new method for solving FDEs. Comparisons are made between the AGM and previously published papers. Several non-linear FDE examples have been solved to illustrate the high performance and quality of the proposed method. Findings show that the AGM is a highly useful and powerful method that can be utilized for FDEs.

© 2024 University of Mazandaran

*Corresponding Author: Khashayar_h68@yahoo.com

Supplementary information: Supplementary information for this article is available at <https://cste.journals.umz.ac.ir/>

Please cite this paper as: Khalili, Z., Bararpour, A., Hosseinzadeh, K., Jafari, B., & Ganji, D. D. (2024). New Approach of Non-Linear Fractional Differential Equations Analytical Solution by Akbari-Ganji's Method. Contributions of Science and Technology for Engineering, 1(1), 12-18.

1. Introduction

FDEs can be used for mathematical modeling of many natural phenomena. These equations are widely used in physics, chemistry, biology, and engineering. Hence, studies have been conducted on the methods of solving these equations. Since there is no exact solution to FDEs, the researchers used various approximate methods to solve this type of equation. To solve FDEs, Sinan et al. [1] employed the Homotopy Perturbation Method (HPM) and found that this method effectively solves higher-order non-linear partial differential equations (PDE). To solve fractional Volterra-Fredholm integro-differential equations, Das et al. [2] employed the HPM, concluding that taking this method may lead to the model's exact solution. The HPM was investigated by Javeed et al. [3] for the solution of fractional partial differential equations (FPDEs). For non-linear FPDEs, Aguilar et al. [4] employed the Homotopy Perturbation Transform Method (HPTM). This method has also been used to solve equations in other studies [5 and 6]. Another method for solving FDEs is the Variational Iteration Method (VIM). DjelloulZiane et al. [7] employed a combination of the VIM and the Laplace transform method to solve linear and non-linear PDEs with time-fractional derivatives; they claimed that the suggested method is appropriate for such issues and that it is highly

efficient. Sontakke et al. [8] employed the VIM to solve time FPDEs and concluded that the VIM is a useful method for solving non-linear FPDEs [9 and 10]. Demir et al. [11] used the differential transform method (DTM) to estimate solutions of fractional Bratu-type differential equations. Their findings show that the DTM can be used for non-linear FDEs in a simple and effective manner, concluding that the current method produces good results. Bansal et al. [12] employed the extended DTM to find analytical exact solutions to two key specific examples of the Bagley Torvik equation, both having Caputo fractional derivatives. Their research shows how the DTM can be used to solve FDEs [13 and 14]. Panda et al. [15] used an Adomian decomposition method (ADM) and HPM to solve a time-fractional partial integrodifferential equation. Abdollah Sadeghinia et al. [16] used the ADM to solve non-linear multi-term FDEs. Haq et al. [17] and Verma et al. [18] conducted more research on the Adomian decomposition method. Other methods, such as the Shehu method [19], the Adams Bashforth-Moulton method [20], the fractional residual power series method (FRPSM) [21], the B-SPLINE OPERATIONAL MATRIX method [22], the Bernoulli polynomials method [23], and Lubich's fractional linear multi-step methods [24], have been proposed in other studies. This form of FDEs [25] is investigated in this paper.






Figure 1. The results of Example 4.1 when $\alpha=1.5$, compared with the results obtained by VIM




Figure 2. The results of Example 4.1 when $\alpha = 0.25, 0.5, 0.75, 1.25$ and 1.75

4.2. Example 2

Consider the following non-linear FDE [25].

$$D^\alpha y(t) + e^t y^3(t) = e^t + e^{4t}, \quad 0 < \alpha \leq 2 \tag{24}$$

According to the boundary conditions:

$$[y(t)]_{t=0} = 1, \quad [y(t)]_{t=1} = e \tag{25}$$

In AGM, the answer of the differential equations is considered as a finite series with constant coefficients as follows:

$$y(t) = \sum_{i=0}^1 a_i t^i = a_0 + a_1 t \tag{26}$$

$$y(\tau) = \sum_{i=0}^3 b_i \tau^i = b_0 + b_1 \tau + b_2 \tau^2 + b_3 \tau^3 \tag{27}$$

Substituting Equations 26 and 27 in Equation 24 as follows:

$$y(t) = 4.513516668 * t^{3/2} * b_3 + 2.256758334 * \sqrt{t} * b_2 + e^t * (t * a_1 + a_0)^3 - e^t - e^{4t} \tag{28}$$

Then, the boundary conditions will be applied as follows:

$$y(0) = 1 \rightarrow a_0 = 1 \tag{29}$$

$$y(1) = 1 \rightarrow a_1 + a_0 = e \tag{30}$$

$$f(y(1)) = 4.513516668 * b_3 + 2.256758334 * b_2 + e * (a_1 + a_0)^3 - e - e^4 = 0 \tag{31}$$

$$f'(y(1)) = 6.770275002 * b_3 + 1.128379167 * b_2 + e * (a_1 + a_0)^3 + 3 * e * (a_1 + a_0)^2 * a_1 - e - 4 * e^4 = 0 \tag{32}$$

The unknown coefficients of Equation 28 will be gained by solving these four equations:

$$\{a_0 = 1.000000000, a_1 = 1.718281828, b_2 = -26.09826182, b_3 = 13.65138455\} \tag{33}$$

By substituting the calculated coefficients in Equation 26, the differential equation will be obtained.

$$y(t) = 1.718281828 * t + 1.000000000 \tag{34}$$

The comparison between AGM results and the results obtained by Nagdy et al. [25] when $\alpha=1.5$ has been shown in Table 2 and Figure 3. As can be seen, the results obtained by AGM are in good agreement with the results obtained by Nagdy et al. [25].

Table 2. Comparison of the results obtained by AGM and VIM at $\alpha = 1.5$

t	Present work	Nagdy et al.	Error (%)
0	1	1	0
0.1	1.171828	1.157112	1.271768
0.2	1.343656	1.322714	1.583271
0.3	1.515485	1.491825	1.585917
0.4	1.687313	1.661503	1.553402
0.5	1.859141	1.829563	1.616645
0.6	2.030969	1.994985	1.803733
0.7	2.202797	2.158847	2.035824
0.8	2.374625	2.325785	2.099946
0.9	2.546454	2.50605	1.612233
1	2.718282	2.7183	0.000669

The results of Example 4.2 when $\alpha = 0.25, 0.5, 0.75, 1.25,$ and 1.75 are shown in Figure 4. As shown in Figure 4, by decreasing $\alpha, y(t)$ has been increased.




Figure 3. The results of Example 4.2 when $\alpha=1.5$, compared with the results obtained by VIM




Figure 4. The results of Example 4.2 when $\alpha = 0.25, 0.5, 0.75, 1.25, \text{ and } 1.75$

4.3. Example 3

Consider the following non-linear FDE [25].

$$D^\alpha y(t) + ty^2(t) = -sint + t\sin^2(t), \quad 0 < \alpha \leq 2, \quad (35)$$

According to the boundary conditions:

$$[y(t)]_{t=0} = 0, \quad [y(t)]_{t=\frac{\pi}{2}} = 1 \quad (36)$$

In AGM, the answer of the differential equations is considered as a finite series with constant coefficients as follows:

$$y(t) = \sum_{i=0}^3 a_i t^i = a_0 + a_1 t + a_2 t^2 + a_3 t^3 \quad (37)$$

$$y(\tau) = \sum_{i=0}^3 b_i \tau^i = b_0 + b_1 \tau + b_2 \tau^2 + b_3 \tau^3 \quad (38)$$

Substituting Equations 37 and 38 in Equation 35 as follows:

$$y(t) = 4.513516668 * t^{3/2} * b_3 + 2.256758334 * \sqrt{t} * b_2 + t * (t^3 * a_3 + t^2 * a_2 + t * a_1 + a_0)^2 + \sin(t) - t * \sin(t)^2 = 0 \quad (39)$$

Then, the boundary conditions will be applied as follows:

$$y(0) = 0 \rightarrow a_0 = 0 \quad (40)$$

$$y\left(\frac{\pi}{2}\right) = \frac{1}{8} * \pi^3 * a_3 + \frac{1}{4} * \pi^2 * a_2 + \frac{1}{2} * \pi * a_1 + a_0 = 1 \quad (41)$$

$$f(y(1)) = \left(1.128379167 * \sqrt{2} * \pi^2\right) * b_3 + 1.128379167 * \sqrt{2} * \sqrt{\pi} * b_2 + \frac{1}{2} * \pi * \left(\frac{1}{8} * \pi^3 * a_3 + \frac{1}{4} * \pi^2 * a_2 + \frac{1}{2} * \pi * a_1 + a_0\right)^2 + 1 - \frac{1}{2} * \pi = 0 \quad (42)$$

$$f'(y(1)) = 3.385137501 * \sqrt{2} * \sqrt{\pi} * b_3 + \frac{1.128379167 * \sqrt{2} * b_2}{\sqrt{\pi}} + \left(\frac{1}{8} * \pi^3 * a_3 + \frac{1}{4} * \pi^2 * a_2 + \frac{1}{2} * \pi * a_1 + a_0\right) * \left(\frac{3}{4} * \pi^2 * a_3 + \pi * a_2 + a_1\right) - 1 = 0 \quad (43)$$

$$f''(y(1)) = \frac{3.385137501 * \sqrt{2} * b_3}{\sqrt{\pi}} - \frac{1.128379167 * \sqrt{2} * b_2}{\sqrt{\pi^{3/2}}} + 4 * \left(\frac{1}{8} * \pi^3 * a_3 + \frac{1}{4} * \pi^2 * a_2 + \frac{1}{2} * \pi * a_1 + a_0\right) * \left(\frac{3}{4} * \pi^2 * a_3 + \pi * a_2 + a_1\right) + \pi * \left(\frac{3}{4} * \pi^2 * a_3 + \pi * a_2 + a_1\right)^2 + \pi * \left(\frac{1}{8} * \pi^3 * a_3 + \frac{1}{4} * \pi^2 * a_2 + \frac{1}{2} * \pi * a_1 + a_0\right) * (3 * \pi * a_3 + 2 * a_2) - 1 + \pi = 0 \quad (44)$$

$$f'(y(1)) = \frac{3.385137501 * \sqrt{2} * b_3}{\sqrt{\pi^{3/2}}} + \frac{3.385137501 * \sqrt{2} * b_2}{\sqrt{\pi^{5/2}}} + 6 * \left(\frac{3}{4} * \pi^2 * a_3 + \pi * a_2 + a_1\right)^2 + 6 * \left(\frac{1}{8} * \pi^3 * a_3 + \frac{1}{4} * \pi^2 * a_2 + \frac{1}{2} * \pi * a_1 + a_0\right) * (3 * \pi * a_3 + 2 * a_2) + 3 * \pi * \left(\frac{3}{4} * \pi^2 * a_3 + \pi * a_2 + a_1\right) * (3 * \pi * a_3 + 2 * a_2) + 6 * \pi * \left(\frac{1}{8} * \pi^3 * a_3 + \frac{1}{4} * \pi^2 * a_2 + \frac{1}{2} * \pi * a_1 + a_0\right) * a_3 + 6 = 0 \quad (45)$$

The unknown coefficients of Equation 39 will be gained by solving these six equations.

$$\begin{aligned} a_0 &= 0, \\ a_1 &= 1.147465869, \\ a_2 &= -0.2828944409, \\ a_3 &= -0.02694193005, \\ b_2 &= -0.4268734354, \\ b_3 &= 0.02333849508 \end{aligned} \quad (46)$$

By substituting the calculated coefficients in Equation 37, the differential equation will be obtained.

$$y(t) = -0.02694193005 * t^3 - 0.2828944409 * t^2 + 1.147465869 * t \quad (47)$$

The comparison between AGM results and the results obtained by Nagdy et al. [25] when $\alpha=1.5$ has been shown in Table 3 and Figure 5. As can be seen, the results obtained by AGM are in good agreement with the results obtained by Nagdy et al. [25].

Table 3. Comparison of the results obtained by AGM and VIM at $\alpha = 1.5$

t	Present work	Nagdy et al.	Error (%)
0	0	0	0
0.2	0.214194	0.190696	12.32212
0.4	0.405351	0.372123	8.929335
0.6	0.572248	0.540419	5.889781
0.8	0.713661	0.692453	3.062817
1	0.828367	0.824106	0.517031
1.2	0.915141	0.928384	1.42648
1.4	0.97276	0.993341	2.071898
1.6	1	1.000094	0.009422



Figure 5. The results of Example 4.3 when $\alpha=1.5$, compared with the results obtained by VIM

The results of Example 4.3 when $\alpha = 0.25, 0.5, 0.75, 1.25, 1.5$ and 1.75 are shown in Figure 6.




Figure 6. The results of Example 4.3 when $\alpha = 0.25, 0.5, 0.75, 1.25$ and 1.75

As shown in Figure 6, by decreasing α , $y(t)$ has been increased.

5. Conclusion

The AGM was successfully employed to find a solution to non-linear FDEs in this study. Three examples have been chosen to demonstrate the method's efficiency and accuracy. The VIM was compared with the approximate answers generated by the AGM, and the results of the equations are shown in Tables 1 to 3 to demonstrate the method's stability. Then, the equations were solved for various fractional orders, and the results show that as the α decreases, the value of $y(t)$ increases. Results revealed that AGM is useful and powerful for solving FDEs.

6. References

- [1] Sinan, M., Shah, K., Khan, Z. A., Al-Mdallal, Q., & Rihan, F. (2021). On semianalytical study of fractional-order kawahara partial differential equation with the homotopy perturbation method. *Journal of Mathematics*, 2021. doi:10.1155/2021/6045722.
- [2] Das, P., Rana, S., & Ramos, H. (2019). Homotopy perturbation method for solving Caputo- type fractional-order Volterra- Fredholm integro- differential equations. *Computational and Mathematical Methods*, 1(5). doi:10.1002/cmm4.1047.
- [3] Javeed, S., Baleanu, D., Waheed, A., Shaukat Khan, M., & Affan, H. (2019). Analysis of Homotopy Perturbation Method for Solving Fractional Order Differential Equations. *Mathematics*, 7(1), 40. doi:10.3390/math7010040.
- [4] Gómez-Aguilar, J., Yépez-Martínez, H., Torres-Jiménez, J., Córdova-Fraga, T., Escobar-Jiménez, R., & Olivares-Peregrino, V. (2017). Homotopy perturbation transform method for nonlinear differential equations involving to fractional operator with exponential kernel. *Advances in Difference Equations*, 2017(1). doi:10.1186/s13662-017-1120-7.
- [5] Inas, A. A., & Eladdad, E. E. (2022). On Solving Partial Differential Equations by a Coupling of the Homotopy Perturbation Method and a New Integral Transform. *Applied Mathematics and Information Sciences*, 16(1), 51–57. doi:10.18576/amis/160106.
- [6] Shabeeb, M. A., Aljaboori, M. A., & Jasim, S. H. (2018). Homotopy Perturbation Method For Solving Coupled Of Partial Differential Equation. *مجلة ميسان للدراسات الأكاديمية*, ٣٢٦. doi:10.54633/2333-017-034-029.
- [7] Ziane, D., & Cherif, M. H. (2018). Variational iteration transform method for fractional differential equations. *Journal of Interdisciplinary Mathematics*, 21(1), 185–199. doi:10.1080/09720502.2015.1103001.
- [8] Sontakke, B. R., Shelke, A. S., & Shaikh, A. S. (2019). Solution of Non-Linear Fractional Differential Equations By Variational Iteration Method and Applications. *Far East Journal of Mathematical Sciences (FJMS)*, 110(1), 113–129. doi:10.17654/ms110010113.
- [9] Singh, B. K., & Kumar, P. (2017). Fractional Variational Iteration Method for Solving Fractional Partial Differential

- Equations with Proportional Delay. *International Journal of Differential Equations*, 2017. doi:10.1155/2017/5206380.
- [10] Sakar, M. G., & Ergören, H. (2015). Alternative variational iteration method for solving the time-fractional Fornberg-Whitham equation. *Applied Mathematical Modelling*, 39(14), 3972–3979. doi:10.1016/j.apm.2014.11.048.
- [11] Demir, D. D., & Zeybek, A. (2017). The Numerical Solution of Fractional Bratu-Type Differential Equations. *ITM Web of Conferences*, 13, 01008. doi:10.1051/itmconf/20171301008.
- [12] Bansal, M. K., & Jain, R. (2016). Analytical solution of bagley Torvik equation by generalize differential transform. *International Journal of Pure and Applied Mathematics*, 110(2), 265–273. doi:10.12732/ijpam.v110i2.3.
- [13] Ünal, E., & Gökdoğan, A. (2017). Solution of conformable fractional ordinary differential equations via differential transform method. *Optik*, 128, 264–273. doi:10.1016/j.ijleo.2016.10.031.
- [14] Yang, X. J., Tenreiro Machado, J. A., & Srivastava, H. M. (2016). A new numerical technique for solving the local fractional diffusion equation: Two-dimensional extended differential transform approach. *Applied Mathematics and Computation*, 274, 143–151. doi:10.1016/j.amc.2015.10.072.
- [15] Panda, A., Santra, S., & Mohapatra, J. (2022). Adomian decomposition and homotopy perturbation method for the solution of time fractional partial integro-differential equations. *Journal of Applied Mathematics and Computing*, 68(3), 2065–2082. doi:10.1007/s12190-021-01613-x.
- [16] Sadeghinia, A., & Kumar, P. (2021). One Solution of Multi-term Fractional Differential Equations by Adomian Decomposition Method: Scientific Explanation. *Current Topics on Mathematics and Computer Science Vol. 6*, 120–130. doi:10.9734/bpi/ctmcs/v6/11542d.
- [17] Haq, F., Shah, K., ur Rahman, G., & Shahzad, M. (2018). Numerical solution of fractional order smoking model via laplace Adomian decomposition method. *Alexandria Engineering Journal*, 57(2), 1061–1069. doi:10.1016/j.aej.2017.02.015.
- [18] Verma, P., & Kumar, M. (2022). Analytical solution with existence and uniqueness conditions of non-linear initial value multi-order fractional differential equations using Caputo derivative. *Engineering with Computers*, 38(1), 661–678. doi:10.1007/s00366-020-01061-4.
- [19] Khalouta, A., & Kadem, A. (2019). A New Method to Solve Fractional Differential Equations: Inverse Fractional Shehu Transform Method. *An International Journal (AAM)*, 14(2), 926–941. <http://pvamu.edu/aam>
- [20] Baskonus, H. M., & Bulut, H. (2015). On the numerical solutions of some fractional ordinary differential equations by fractional Adams-Bashforth-Moulton method. *Open Mathematics*, 13(1), 547–556. doi:10.1515/math-2015-0052.
- [21] Khalouta, A., & Kadem, A. (2020). An efficient method for solving nonlinear time-fractional wave-like equations with variable coefficients. *Tbilisi Mathematical Journal*, 12(4). doi:10.32513/tbilisi/1578020573.
- [22] Jafari, H., Tajadodi, H., & Baleanu, D. (2015). A numerical approach for fractional order riccati differential equation using B-spline operational matrix. *Fractional Calculus and Applied Analysis*, 18(2), 387–399. doi:10.1515/fca-2015-0025.
- [23] Keshavarz, E., Ordokhani, Y., & Razzaghi, M. (2016). A numerical solution for fractional optimal control problems via Bernoulli polynomials. *JVC/Journal of Vibration and Control*, 22(18), 3889–3903. doi:10.1177/1077546314567181.
- [24] Garrappa, R. (2018). Numerical Solution of Fractional Differential Equations: A Survey and a Software Tutorial. *Mathematics*, 6(2), 16. doi:10.3390/math6020016.
- [25] Nagdy, A. S., & Hashem, K. M. (2020). Numerical solutions of nonlinear fractional differential equations by variational iteration method. *Journal of Nonlinear Sciences and Applications*, 14(02), 54–62. doi:10.22436/jnsa.014.02.01.
- [26] Meng, Z., Yi, M., Huang, J., & Song, L. (2018). Numerical solutions of nonlinear fractional differential equations by alternative Legendre polynomials. *Applied Mathematics and Computation*, 336, 454–464. doi:10.1016/j.amc.2018.04.072.
- [27] Loverro, A. (2004). Fractional calculus: history, definitions and applications for the engineer. *Rapport technique*, Department of Aerospace and Mechanical Engineering, Univeristy of Notre Dame, Notre Dame, United States.
- [28] Akbari, M. R., Ganji, D. D., Nimafar, M., & Ahmadi, A. R. (2014). Significant progress in solution of nonlinear equations at displacement of structure and heat transfer extended surface by new AGM approach. *Frontiers of Mechanical Engineering*, 9(4), 390–401. doi:10.1007/s11465-014-0313-y.